

Atmospheric circulation changes identified thanks to ground-based FTIR monitoring of hydrogen chloride (HCl)

Emmanuel Mahieu, University of Liège, Liège, Belgium

Martyn Chipperfield, University of Leeds, Leeds, U.K.

Justus Notholt, University of Bremen, Bremen, Germany

Thomas Reddmann, Karlsruhe Institute of Technology, Karlsruhe, Germany

Monitoring the success of the Montreal Protocol on substances that deplete stratospheric ozone is one of the primary tasks of the NDACC network. Among the various techniques involved, giving access to numerous relevant parameters, the ground-based FTIR instruments contribute significantly by providing total and partial columns of key tropospheric and stratospheric constituents. Indeed, high-resolution solar infrared spectra contain the signatures of a suite of halogenated organic source gases. The current list includes CFC-11, CFC-12, HCFC-22, HCFC-142b, CCl_4 , CF_4 , SF_6 (e.g., Krieg et al., 2005; Zander et al., 2008; Rinsland et al., 2012; Mahieu et al., 2013a; Mahieu et al., 2014a) and efforts are ongoing to expand this list. In addition, the respective evolutions of the inorganic chlorine and fluorine loadings in the stratosphere are also accessible to this technique through observations of the main reservoirs, i.e. hydrogen chloride (HCl) and chlorine nitrate (ClONO_2), hydrogen fluoride (HF) and carbonyl fluoride (COF_2). Time series and trends of all these species have been reported and analyzed in successive studies, notably allowing characterizing from the ground the rapid increase of inorganic chlorine (Cl_y) in the Earth's stratosphere (e.g., Zander et al., 1987; Rinsland et al., 1991, 1996; Reisinger et

al., 1995), following large emissions of anthropogenic halogenated source gases during the 1970s to 1990s. Later, studies involving several NDACC ground-based FTIR stations provided evidence for a stabilization of HCl and ClONO_2 around the mid-1990s (Rinsland et al., 2003), and then a near-global characterization of the Cl_y decrease at rates close to 1%/yr in both hemispheres at 17 sites between 80°N and 78°S (Kohlhepp et al., 2012).


Recently, Mahieu et al. (2013b) concentrated on the post-peak evolution of HCl using the latest observations available from the Jungfraujoch station (Swiss Alps, 46.5°N) and from a composite satellite partial column time series based on infrared solar occultation measurements performed by HALOE (Halogen Occultation Experiment, onboard UARS; Russell et al., 1993) until 2005 and by ACE-FTS (Atmospheric Chemistry Experiment-Fourier Transform Spectrometer, onboard SCISAT; Bernath et al., 2005) from 2004 onwards. Although this study confirmed for the 1997-2007 time period the amplitude of the downward trend determined by Kohlhepp et al. (2012), both the ground-based and satellite data sets revealed a significant re-increase (at the 2σ level of uncertainty) of the HCl reservoir in the Northern Hemisphere mid-latitude stratosphere after 2007. This feature was unexpected given the well documented and smooth decrease of total organic chlorine in the troposphere (e.g. WMO 2010, 2011). At that time it was possible to exclude that a change in the partitioning among the main inorganic chlorine reservoirs was responsible for the HCl upturn since the Jungfraujoch time series combining HCl and ClONO_2 columns also exhibited a significant increase of Cl_y after 2007. Possible causes for this HCl upturn include a change in atmospheric circulation or the substantial contribution of new unknown chlorine-bearing source gases to the organic chlorine budget monitored by the AGAGE and NOAA/ESRL in situ networks.

These observations and questions stimulated a follow up study

including data from other NDACC stations to check whether the recent HCl increase was a global atmospheric feature. We also used the GOZCARDS satellite data set (merging measurements of HALOE, ACE-FTS and Aura/MLS; Froidevaux et al., 2013) in order to confirm the ground-based observations and to get information

on the altitude range where the changes were taking place as well as simulations by the state-of-the-art 3-D chemistry transport models SLIMCAT and KASIMA (Kohlhepp et al., 2012).

The paper reporting about these investigations has recently been published in Nature (Mahieu et al., 2014b) and the main findings






Figure 2. Mean differences of the HCl concentration [panel a) and age-of-air (panel b) between 2010/2011 and 2005/2006 as a function of altitude and latitude, as derived from the standard SLIMCAT run [from Mahieu et al., doi:10.1038/nature13857, 2014b).

are as follows:

- As shown in Figure 1a, HCl negative rates of changes are found at the eight NDACC sites for 1997-2007 (in red), in agreement with the satellite data (in orange); SLIMCAT and KASIMA model simulations using ERA-Interim meteorology and surface source gases mixing ratios from the WMO A1 scenario confirm these trends in sign and amplitude (in green and grey, respectively).

- For 2007-2011, we observed a contrasting situation, with NDACC, GOZCARDS, SLIMCAT and KASIMA showing significant increases of HCl in the Northern Hemisphere, at rates up to 3%/yr, while the various data sets consistently show HCl decreases in the Southern Hemisphere.

- A dedicated SLIMCAT run (denoted S2000, light green) used constant 6-hourly winds of 2000, from 2000 onwards, to study the impact of atmospheric dynamics. Clearly, this run does not produce the HCl increase in the Northern Hemisphere (Fig. 1b) and since the only difference between the standard and S2000 SLIMCAT run resides in the meteorological forcings, we conclude that these are responsible for the HCl upturn.

Stratospheric circulation changes have been evaluated using an indicator of mean age-of-air available in both models as an idealized tracer with a linearly increasing tropospheric mixing ratio. Figure 2 shows the spatial distribution of the HCl concentration and mean age-of-air changes between 2005/2006 and 2010/2011, as derived from the standard SLIMCAT run. These plots indicate an obvious asymmetry between both hemispheres, with a clear correlation in the lower stratosphere between the HCl increase/decrease over the time period under investigation and a slower/faster Brewer-Dobson circulation.

The slower and specific transport pathways followed by the air masses over the recent years in the Northern Hemisphere has allowed a longer exposure to strong ultraviolet radiation responsible of the photolysis of the source gases, ultimately resulting in a larger relative conversion into the reservoir species.

The most important point is that by combining the NDACC observations with state-of-the-art CTM simulations, we

can conclude that the recent HCl increase which occurred in the Northern Hemisphere over 2007-2011 does not result from rogue emissions of chlorinated source gases. Hence, the Montreal Protocol is well on track and will lead to an overall reduction of the stratospheric chlorine abundance required to ensure ozone recovery in the decades to come. Our study also underlines the constant need for reliable long-term monitoring of the stratospheric composition, both from the ground and from space.

References

- Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P.-F., Colin, R., DeCola, P., De Mazière, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn, E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nchititu, F., Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelton, R., Sloan, J. J., Soucy, M.-A., Strong, K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R. and Zou, J.: Atmospheric Chemistry Experiment (ACE): Mission overview, *Geophysical Research Letters*, 32(15), doi:10.1029/2005GL022386, 2005.
- Froidevaux, L., Anderson, J., Fuller, R. A., Bernath, P. F., Livesey, N. J., Russell III, J. M., and Walker, K. A.: GOZCATS Merged Data for Hydrogen Chloride Monthly Zonal Means on a Geodetic Latitude and Pressure Grid version 1.01, <http://dx.doi.org/10.5067/MEASURES/GOZCATS/DATA3002>, NASA Goddard Earth Science Data and Information Services Center, accessed June 2013.
- Kohlhepp, R., Ruhnke, R., Chipperfield, M. P., De Mazière, M., Notholt, J., Barthlott, S., Batchelor, R. L., Blatherwick, R. D., Blumenstock, T., Coffey, M. T., Demoulin, P., Fast, H., Feng, W., Goldman, A., Griffith, D. W. T., Hamann, K., Hannigan, J. W., Hase, F., Jones, N. B., Kagawa, A., Kaiser, I., Kasai, Y., Kirner, O., Kouker, W., Lindenmaier, R., Mahieu, E., Mittermeier, R. L., Monge-Sanz, B., Morino, I., Murata, I., Nakajima, H., Palm, M., Paton-Walsh, C., Raffalski, U., Reddman, T., Rettinger, M., Rinsland, C. P., Rozanov, E., Schneider, M., Senten, C., Servais, C., Sinnhuber, B.-M., Smale, D., Strong, K., Sussmann, R., Taylor, J. R., Vanhaelewyn, G., Warneke, T., Whaley, C., Wiehle, M. and Wood, S. W.: Observed and simulated time evolution of HCl, ClONO₂, and HF total column abundances, *Atmospheric Chemistry and Physics*, 12(7), 3527–3556, doi:10.5194/acp-12-3527-2012, 2012.
- Krieg, J., Notholt, J., Mahieu, E., Rinsland, C. P. and Zander, R.: Sulphur hexafluoride (SF₆): comparison of FTIR-measurements at three sites and determination of its trend in the northern hemisphere, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 92(3), 383–392, doi:10.1016/j.jqsrt.2004.08.005, 2005.
- Mahieu, E., O'Doherty, S., Reimann, S., Vollmer, M. K., Bader, W., Bovy, B., Lejeune, B., Demoulin, P., Roland, G., Servais, C. and Zander, R.: First retrievals of HCFC-142b from ground-based high-resolution FTIR solar observations: application to high-altitude Jungfraujoch spectra, *Geophysical Research Abstracts*, 15, EGU2013-1185-1, 2013a.
- Mahieu, E., Zander, R., Bernath, P. F., Boone, C. D. and Walker, K. A.: Recent trend anomaly of hydrogen chloride (HCl) at northern mid-latitudes derived from Jungfraujoch, HALOE and ACE-FTS Infrared solar observations, in *The Atmospheric Chemistry Experiment ACE at 10: A Solar Occultation Anthology*, P.F. Bernath (Ed.), ISBN 978-0-937194-54-9, A. Deepak Publishing, 239-249, 2013b.
- Mahieu, E., Zander, R., Toon, G. C., Vollmer, M. K., Reimann, S., Mühle, J., Bader, W., Bovy, B., Lejeune, B., Servais, C., Demoulin, P., Roland, G., Bernath, P. F., Boone, C. D., Walker, K. A. and Duchatelet, P.: Spectrometric monitoring of atmospheric carbon tetrafluoride (CF₄) above the Jungfraujoch station since 1989: evidence of continued increase but at a slowing rate, *Atmospheric Measurement Techniques*, 7(1), 333–344, doi:10.5194/amt-7-333-2014, 2014a.
- Mahieu, E., Chipperfield, M. P., Notholt, J., Reddman, T., Anderson, J., Bernath, P. F., Blumenstock, T., Coffey, M. T., Dhomse, S. S., Feng, W., Franco, B., Froidevaux, L., Griffith, D. W. T., Hannigan, J. W., Hase, F., Hossaini, R., Jones, N. B., Morino, I., Murata, I., Nakajima, H., Palm, M., Paton-Walsh, C., Russell III, J. M., Schneider, M., Servais, C., Smale, D. and Walker, K. A.: Recent Northern Hemisphere stratospheric HCl increase due to atmospheric circulation changes, *Nature*, 515 (7525), 104-107, doi:10.1038/nature13857, 2014b.
- Reisinger, A. R., Jones, N. B., Matthews, W. A. and Rinsland, C. P.: Southern hemisphere midlatitude ground-based measurements of ClONO₂: Method of analysis, seasonal cycle and long term-trend, *Journal of Geophysical Research*, 100(D11), 23183, doi:10.1029/95JD02529, 1995.
- Rinsland, C. P., Levine, J. S., Goldman, A., Sze, N. D., Ko, M. K. W. and Johnson, D. W.: Infrared measurements of HF and HCl total column abundances above Kitt Peak, 1977–1990: Seasonal cycles, long-term increases, and comparisons with model calculations, *Journal of Geophysical Research*, 96(D8), 15523-15540, doi:10.1029/91JD01249, 1991.
- Rinsland, C. P., Zander, R., Demoulin, P. and Mahieu, E.: ClONO₂ total vertical column abundances above the Jungfraujoch Station, 1986–1994: Long-term trend and winter-spring enhancements, *Journal of Geophysical Research*, 101(D2), 3891, doi:10.1029/95JD03349, 1996.
- Rinsland, C. P., Mahieu, E., Zander, R., Jones, N. B., Chipperfield, M. P., Goldman, A., Anderson, J., Russell III, J. M., Demoulin, P., Notholt, J., Toon, G. C., Blavier, J.-F., Sen, B., Sussmann, R., Wood, S. W., Meier, A., Griffith, D. W. T., Chiou, L. S., Murracay, F. J., Stephen, T. M., Hase, F., Mikuteit, S., Schulz, A. and Blumenstock, T.: Long-term trends of inorganic chlorine from ground-based infrared solar spectra: Past increases and evidence for stabilization, *Journal of Geophysical Research: Atmospheres*, 108(D8), doi:10.1029/2002JD003001, 2003.
- Rinsland, C. P., Mahieu, E., Demoulin, P., Zander, R., Servais, C. and Hartmann, J.-M.: Decrease of the carbon tetrachloride (CCl₄) loading above Jungfraujoch, based on high resolution infrared solar spectra recorded between 1999 and 2011, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113(11), 1322–1329, doi:10.1016/j.jqsrt.2012.02.016, 2012.
- Russell III, J. M., Gordley, L. L., Park, J. H., Drayson, S. R., Hesketh, W. D., Cicerone, R. J., Tuck, A. F., Frederick, J. E., Harries, J. E. and Crutzen, P. J.: The Halogen Occultation Experiment, *Journal of Geophysical Research*, 98(D6), 10777, doi:10.1029/93JD00799, 1993.
- Zander, R., Roland, G., Delbouille, L., Sauval, A., Farmer, C. B. and Norton, R. H.: Column abundance and the long-term trend of hydrogen chloride (HCl) above the Jungfraujoch Station, *Journal of Atmospheric Chemistry*, 5(4), 395–404, doi:10.1007/BF00113902, 1987.
- Zander, R., Mahieu, E., Demoulin, P., Duchatelet, P., Roland, G., Servais, C., De Mazière, M., Reimann, S. and Rinsland, C. P.: Our changing atmosphere: Evidence based on long-term infrared solar observations at the Jungfraujoch since 1950, *Science of The Total Environment*, 391(2-3), 184–195, doi:10.1016/j.scitotenv.2007.10.018, 2008.
- WMO (World Meteorological Organization): Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project, Report N° 52, Geneva, Switzerland, 2011.